

# NALOXONE

## BE PREPARED



### AT RISK

- HIGHER DOSE PRESCRIPTION OPIOIDS
- OPIOIDS + BENZOS COMBINED
- OPIOID DEPENDENTS
- INJECTING OPIOIDS



### SIGNS OF OVERDOSE

- UNCONSCIOUS OR NON-RESPONSIVE
- BLUE SKIN/LIPS
- PINPOINT PUPILS
- SLOW/NO BREATHING

## HOW TO RESPOND



## CALL 999

### UNCONSCIOUS + BREATHING

- ADMINISTER NALOXONE
- RECOVERY POSITION

### UNCONSCIOUS NOT BREATHING


- ADMINISTER NALOXONE
- START CPR



## NO

- WALKING
- SHOWER/BATH
- SLAPPING ETC

## FOLLOW INSTRUCTIONS <sup>IN</sup> KIT



### INJECTION (PRENOXAD)

- Keep Sealed
- Twist cap off
- Inject into arm or thigh
- One dose every 2-3 mins until responsive
- Place in original box between doses
- Sharps bin after usage



### NASAL SPRAY (NYXOID)

- Do no test beforehand
- Tilt head backwards
- Place spray in nostril, press plunger until clicks
- Another dose every 2-3 mins until responsive



## REPLACE/REFRESH KIT AS NEEDED